



APRIL/MAY 2019

MMA41 — COMPLEX ANALYSIS — II

Time : Three hours

Maximum : 75 marks

SECTION A — (5 × 6 = 30 marks)

Answer ALL questions.

1. (a) Prove that the infinite product  $\prod_1 (1 + \alpha_n)$  with  $1 + \alpha_n \neq 0$  converges simultaneously with the series  $\sum_1 \log(1 + \alpha_n)$  whose terms represent the values of the principal branch of the logarithm.

Or

- (b) Show that  $\prod_{n=2}^{\infty} \left(1 - \frac{1}{n^2}\right) = \frac{1}{2}$ .

2. (a) Prove that for  $\sigma = \text{Res} > 1$ ,

$$\frac{1}{\zeta(s)} = \prod_{n=1}^{\infty} (1 - p_n^{-s}).$$

Or



- (b) Prove that a family  $F$  is normal if and only if its closure  $F^-$  with respect to the distance

function  $\rho(f, g) = \sum_{k=1}^{\infty} \delta_k(f, g) 2^{-k}$  is compact.

3. (a) Prove that a continuous function  $u(z)$  which satisfies condition

$$u(z_0) = \frac{1}{2\pi} \int_0^{2\pi} u(z_0 + re^{i\theta}) d\theta$$

is necessarily harmonic.

Or

- (b) Describe the Harnack's principle.

4. (a) Show that an elliptic function without poles is constant.

Or

- (b) Prove that the zeros  $a_1, \dots, a_n$  and poles  $b_1, \dots, b_n$  of an elliptic function satisfy  $a_1 + \dots + a_n \equiv b_1 + \dots + b_n \pmod{M}$ .

5. (a) Prove that the modular function  $\lambda(\tau)$  effects a one-to-one conformal mapping of the region  $\Omega$  onto the upper half plane. The mapping extends continuously to the boundary in such a way that  $\tau = 0, 1, \infty$  correspond to  $\lambda = 1, \infty, 0$ .

Or

2



- (b) Show that any elliptic function with periods

$\omega_1, \omega_2$  can be written as  $C \prod_{k=1}^n \frac{\sigma(z - a_k)}{\sigma(z - b_k)}$

( $C = \text{const.}$ )

### SECTION B — (3 × 15 = 45 marks)

Answer any THREE questions.

6. Derive Jensen's formula.
7. Prove that  $\zeta(s) = 2^s \pi^{s-1} \sin \frac{\pi s}{2} \Gamma(1-s) \zeta(1-s)$ .
8. Prove that given any simply connected region  $\Omega$  which is not the whole plane, and a point  $z_0 \in \Omega$ , there exists a unique analytic function  $f(z)$  in  $\Omega$ , normalized by the conditions  $f(z_0) = 0$ ,  $f'(z_0) > 0$ , such that  $f(z)$  defines a one-to-one mapping of  $\Omega$  onto the disk  $|w| < 1$ .
9. Prove that there exists a basis  $(\omega_1, \omega_2)$  such that the ratio  $\tau = \omega_2 / \omega_1$  satisfies the following conditions: (a)  $\text{Im } \tau > 0$ , (b)  $-\frac{1}{2} < \text{Re } \tau \leq \frac{1}{2}$ , (c)  $|\tau| \geq 1$ , (d)  $\text{Re } \tau \geq 0$  if  $|\tau| = 1$ . The ratio  $\tau$  is uniquely determined by these conditions, and there is a choice of two, four, or six corresponding bases.
10. Derive the Legendre's relation.